

FOR ANIMAL USE ONLY

Enterotect P

FOR SHEEP & GOATS

Reg. No. G4256 (Act 36/1947)

INDICATIONS

Enterotect P is used for vaccinating healthy sheep and goats as an aid in preventing pneumonic pasteurellosis caused by *Mannheimia haemolytica*, enterotoxaemia caused by *Clostridium perfringens* Type A and pulpy kidney caused by *Clostridium perfringens* Type D.

Enterotect P is safe for use in pregnant animals and maternal antibodies are transferred to newborns through the intake of immune colostrum within 48 hours after birth.

COMPOSITION

Alum-precipitated suspensions of *M. haemolytica* leukotoxin with inactivated *C. perfringens* Type A (alpha toxoid) and *C. perfringens* Type D (epsilon toxoid) for use in sheep and goats.

STORAGE

- Store in a refrigerator between 2 °C to 8 °C.
- Do not freeze.
- Protect from high temperatures and sunlight during vaccination.
- Do not use **Enterotect P** after the expiry date printed on the container.

WARNINGS

- **Withdrawal period:** Animals should not be slaughtered for human consumption within 21 days of vaccination.
- Vaccinate healthy animals only.
- Use the entire contents of the vial once opened.
- Discard any empty containers and any unused contents in accordance with local waste disposal regulations.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE

Use only as directed. Shake gently and thoroughly by end over end mixing before and during use. Use sterile needles and syringes only. Do not use chemical sterilisation as traces of disinfectant may negatively affect the vaccine. Do not mix **Enterotect P** with any other vaccines or substances.

DOSAGE

Administer a single 1 ml dose subcutaneously in healthy sheep and goats followed by a second dose 4 to 6 weeks later. The booster vaccination is very important for the development of immunity.

PRESENTATION

Natural white, see-through, high-density polyethylene (HDPE) 20 ml and 100 ml vials closed with gold coloured aluminium combination seals.

REGISTRATION HOLDER & MANUFACTURER

Design Biologix CC

Reg. No. 1992/028856/23

Tel: +27 (0) 12 349 2772

www.designbio.co.za

Enterotect

VIR SKAPE & BOKKE
Reg. Nr. G4256 (Wet 36/1947)

INDIKASIES

Enterotect P word gebruik vir die inenting van gesonde skape en bokke as 'n hulpmiddel in die voorkoming van pneumoniese pasteurellose wat veroorsaak word deur *Mannheimia haemolytica*, enterotoksemie (rooiderm) veroorsaak deur *Clostridium perfringens* Tipe A en bloednier veroorsaak deur *Clostridium perfringens* Tipe D.

Enterotect P is veilig vir gebruik in dragtige diere en moederteenliggaampies word binne 48 uur na geboorte oorgedra aan pasgeborenes deur die inname van kolostrum.

SAMESTELLING

Aluin-gepresipiteerde suspensie van *M. haemolytica* leukotoksien met geïnaktiveerde *C. perfringens* Tipe A (alfa toksoïed) en *C. perfringens* Tipe D (epsilon toksoïed) vir gebruik in skape en bokke.

BERGING

- Berg in 'n yskas tussen 2 °C en 8 °C.
- Moenie vries nie.
- Vermoed blootstelling aan hoë temperature en direkte sonlig gedurende enting.
- Moenie **Enterotect P** gebruik na die vervaldatum wat op die bottel gedruk is nie.

WAARSKUWINGS

- **Onttrekkingsperiode:** Diere moenie binne 21 dae na enting vir menslike verbruik geslag word nie.
- Ent slegs gesonde diere.
- Moenie ongebruikte entstof berg vir latere gebruik nie.
- Vernietig alle leë houers en ongebruikte vloeistof volgens plaaslike afvalbestuursregulasies.
- **HOU BUITE DIE BEREIK VAN KINDERS, ONINGELIGTE PERSONE EN DIERE.**
- Alhoewel hierdie entstof breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui. Skud liggies en meng deeglik voor en tydens gebruik deur die flessie 'n paar keer onderstebo en terug te dop. Gebruik slegs steriele naalde en spuite. Moenie chemiese sterilisasie gebruik nie, aangesien spore van ontsmettingsmiddels die entstof negatief kan beïnvloed. Moenie **Enterotect P** met ander entstowwe of middels meng nie.

DOSIS

Dien 'n enkele 1 ml dosis onderhuid toe in gesonde skape en bokke, gevolg deur 'n tweede dosis 4 tot 6 weke later. Dié skraagingenting is baie belangrik vir die ontwikkeling van immuniteit.

AANBIEDING

Natuurlike wit, deurskynende, hoëdigheidspoliëteleen (HDPE) 20 ml en 100 ml flessies toegemaak met goudkleurige aluminium kombinasie seëls.

REGISTRASIEHOUER & VERVAARDIGER

Design Biologix BK
Reg. Nr. 1992/028856/23
Tel: +27 (0) 12 349 2772
www.designbio.co.za