

## FOR ANIMAL USE ONLY

Consult a veterinarian before use

# Multisomni

Reg. No. G4357 (Act 36/1947)

## INDICATIONS

**Multisomni** is an adjuvanted multi-component vaccine for the active immunisation of healthy cattle against bovine pneumonia caused by *Pasteurella multocida* type A, *P. multocida* type D and *Histophilus somni*.

## DOSAGE FORM

Vaccine suspension for injection with a 1 ml dose via the subcutaneous route. A booster vaccination of 1 ml should be administered 3 to 4 weeks later. Annual revaccination is recommended thereafter.

## STORAGE INSTRUCTIONS

Store in a refrigerator between 2 to 8 °C. Do not freeze. Protect from high temperatures and direct sunlight. Do not use after the expiry date printed on the packaging and container. Do not store any vials into which a needle or other device has been inserted, for future use.

## WARNINGS

- Do not slaughter animals for human consumption within 21 days of vaccination.
- Usually no marked reaction follows vaccination although a transient swelling may appear at the site of inoculation, and some animals may show a moderate rise of temperature for one or two days.
- Do not use in animal species other than those indicated.
- Do not administer to animals in poor or extremely poor body condition.
- KEEP THE PRODUCT OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice; notify the registration holder and the Registrar.

## PRECAUTIONS

- Follow standard sterile procedures during administration of injections.
- Wear protective clothing, masks, gloves, boots, etc. according to hazard standards.
- Avoid contact of the product with skin, eyes and mouth.

- Do not eat, drink or smoke whilst handling the product.
- Do not contaminate rivers, dams or any water sources with containers or waste.
- Dispose of all empty containers and all unused contents in accordance with the local law.
- Do not mix this product with any other vaccines or medicines.

## DIRECTIONS FOR USE

- Shake the bottle well before and during use with gentle end-over-end mixing.
- Inject cattle subcutaneously on the side of the neck. DO NOT deviate from the recommended route and injection site location.
- Recommended dose for **Multisomni** is 1 ml via the subcutaneous route. A booster vaccination of 1 ml should be administered 3 to 4 weeks later. Annual revaccination is recommended thereafter.
- Animals may therefore require a full vaccination program to be completed before the period of highest risk associated with respiratory pneumonia.
- Cattle may be vaccinated from age of weaning and older.
- It is essential to adhere to the vaccination programme to maintain a satisfactory immune response.

## PRESENTATION

**Multisomni**, a light-yellow liquid with an off-white sediment if left to settle, is presented in a pack size of a 20 ml and 100 ml high density polyethylene (HDPE) vial of natural colour, capped with a gold-coloured aluminium cap, labelled and packed in a product-specific outer cardboard carton.

## REGISTRATION HOLDER AND MANUFACTURER

Design Biologix CC  
Reg. No. CK 1992/028856/23  
442 Rigel Avenue South, Erasmusrand, Pretoria, 0181  
Tel: +27 (0) 12 349 2772  
[www.designbio.co.za](http://www.designbio.co.za)

## MARKETED AND DISTRIBUTED BY

Octavoscene (Pty) Ltd  
Tel: +27 (0) 87 237 9995  
[www.octavoscene.co.za](http://www.octavoscene.co.za)



Octavoscene



# Multisomni

Reg. Nr. G4357 (Wet 36/1947)

## AANDUIDINGS

**Multisomni** is 'n multi-komponent entstof met adjuvant vir die aktiewe inenting van gesonde beeste teen respiratoriese longontsteking, veroorsaak deur *Pasteurella multocida* tipe A, *P. multocida* tipe D en *Histophilus somni*.

## ROETE VAN TOEDIENING

Entstof suspensie vir inspuiting met 'n 1 ml dosis via die onderhuidse roete. 'n Skraagdosis van 1 ml moet 3 tot 4 weke later toegedien word. Daar word daarna 'n jaarlikse her-inenting aanbeveel.

## BERGINGSAAWYSINGS

Bêre in 'n yskas tussen 2 tot 8 °C. Moenie vries nie. Beskerm teen hoë temperature en direkte sonlig. Moenie gebruik na die vervaldatum wat op die verpakking en houer gedruk is nie. Moenie houers waarin 'n naald of ander toestel aangebring is, vir toekomstige gebruik bêre nie.

## WAARSKUWINGS

- Moenie diere binne 21 dae na inenting vir menslike verbruik slag nie.
- Normaalweg volg geen merkbare reaksie op inenting nie, alhoewel 'n kortstondige swelling op die plek van inenting kan voorkom, en sommige diere 'n matige styging in temperatuur vir een of twee dae kan toon.
- Moenie in ander spesies anders as die aangeduide spesies gebruik nie.
- Moet nie toegedien word aan diere in 'n swak of uiters swak liggaamskondisie nie.
- HOU DIE PRODUK BUITE BEREIK VAN KINDERS, ONINGELIGTE PERSONE EN DIERE.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid toestande getoets is, mag dit vaal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer en die Registrateur.

## VOORSORGMATREËLS

- Volg standaard steryliese prosedures tydens toediening van inspuitings.
- Dra beskermende klere, maskers, handskoene, stewels, ens. volgens die risikostandaarde.
- Vermoed kontak met die vel, oë en mond.

- Moet nie eet, drink of rook tydens die hantering van die produk nie.
- Moenie riviere, damme of waterbronne met houers of afval besoedel nie.
- Gooi alle leë houers en alle ongebruikte inhoud volgens die plaaslike wetgewing weg.
- Moenie hierdie produk met enige ander entstof of medisyne meng nie.

## GEBRUIKSAANWYSINGS

- Meng die houer goed voor en tydens gebruik.
- Spuit beeste onderhuids aan die kant van die nek. MOENIE van die aanbevole roete en inspuitplek afwyk nie.
- Die aanbevole dosis vir **Multisomni** is 1 ml via die onderhuidse roete. 'n Skraagdosis moet 3 tot 4 weke later toegedien word. Daar word daarna 'n jaarlikse her-inenting aanbeveel.
- Dit word dus vereis dat die volledige inentingsprogram voltooi moet word voor die periode met die grootste risiko wat verband hou met respiratoriese longontsteking.
- Beeste kan ingeënt word vanaf speenouderdom en ouer.
- Dit is noodsaaklik om die inentingsprogram te volg om 'n bevredigende immuunrespons te handhaaf.

## AANBIEDING

**Multisomni**, 'n liggeel vloeistof met 'n wit sediment, word aangebied in 'n pakgrootte van 'n 20 ml en 100 ml hoëdigtheid-poliëtiileen (HDPE) houer van natuurlike kleur, bedek met 'n goudkleurige aluminium doppie, geetiketteer en verpak in 'n produkspesifieke buitenste karton.

## REGISTRASIEHOUER EN VERVAARDIGER

Design Biologix BK  
Reg. Nr. CK 1992/028856/23  
Rigellaan-suid 442, Erasmusrand, Pretoria, 0181  
Tel: +27 (0) 12 349 2772  
www.designbio.co.za

## BEMARK EN VERSPREI DEUR

Octavoscene (Edms) Bpk  
Tel: +27 (0) 87 237 9995  
www.octavoscene.co.za



Octavoscene

